	+ CS151C 2 710N	OF THIS PAGE "						
ECURITY CL	A22161CA - 101		REPORT DOCUM	ENTATION PAGE				
			REPORT DOCUM					
10. REPORT SECURITY CLASSIFICATION Unclassified				1b. RESTRICTIVE MARKINGS None				
26. SECURITY CLASSIFICATION AUTHORITY				J. DISTRIBUTION/AVAILABILITY OF REPORT  Unlimited Approved for public releaso;  distribution unlimited.				
26. DECLASSIFICATION/DOWNGRADING SCHEDULE								
4. PERFORMING ORGANIZATION REPORT NUMBER(S)				5. MONITORING OR	GANIZA	NUMBER(S)		
						AFOSI	R-TR-96	
Unive:	University of Wisconsin  Department of Physics			AFOSR-TR-96  AFOSR -NE  AFOSR -NE				
6c. ADDRESS (City. State and ZIP Code) 1150 University Avenue Madison, WI 53706				AFOSR, 110 Duncan Ave., Suite Dans Bolling AFB, D.C. 20332				
86. NAME OF FUNDING/SPONSORING ORGANIZATION AFOSR—NE				9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER F49620- 93-1-0450				
8c. ADDRES	8c. ADDRESS (City, State and ZIP Code)				10. SOURCE OF FUNDING NOS.			
110 Duncan Ave., Suite B115				PROGRAM ELEMENT NO.	PROJECT NO.	TASK NO.	WORK UNIT	
Bolling AFB, D.C. 20332				61102F	-2301	<del>DS-</del>		
E '	11. TITLE (Include Security Claufication)  Collisions of Atmospheric Gas Molecules				3484	T-S		
12 PERSON	C. Lin		e das nerecures					
13a TYPE OF REPORT 13b. TIME COVERED FROM 8/1/93 TO 7/31/				14. DATE OF REPORT (Yr., Mo., Day)  September 30, 1996  One				
Ł	MENTARY NO							
17.	17. COSATI CODES 18. S			(Continue on reverse if r	secessary and identi	ly by block numbe	r)	
FIELD	GROUP	SUB. GR.						

ISTRACT (Continue on reverse if necessary and identify by block number)

The absolute electron-impact optical emission cross sections of numerous vibrational bands of the second positive band system of the nitrogen molecule have been measured for electron impact energy from threshold up to 600 eV. From the optical data the apparent electron-impact excitation cross sections for five vibrational levels of the  $C^3\Pi_u$  electronic state of the nitrogen molecule have been determined.

We have constructed a new apparatus for measuring the electron-impact optical cross sections for infrared emissions using a Fourier Transform spectrometer for optical detection. This apparatus allows us to study many important infrared emissions which could not be detected previously by our photomultiplier optical detection systems.

	DITC QUALITY INSPECTED 4				
STRIBUTION/AVAILABILITY OF ABSTRACT	21. ABSTRACT SECURITY CLASSIFICATION				
CLASSIFIED/UNLIMITED SAME AS RPT. C DTIC USERS C	Unclassified				
Dr Ralph Kelley	22b. TELEPHONE NUMBER (Include Area Code) 202767-4908	22c. OFFICE SYMBOL			

DD FORM 1473, 83 APR

EDITION OF 1 JAN 73 IS OBSOLETE.

Unclassified

## Final Technical Report (1 August 93 - 31 July 96)

## (AASERT-93) Collisions of Atmospheric Gas Molecules

## AFOSR Grant F49620-93-1-0450

We have conducted a comprehensive study of the electron excitation of the second positive band system of the  $N_2$  molecule as well as the excitation of the  $C^3\Pi_u$  electronic state of the  $N_2$  molecule. We have measured the absolute optical emission cross sections of numerous (v', v'') vibrational bands of the second positive system for electron impact energy from threshold up to 600 eV. The relative optical emission cross sections are compared with theoretical calculations. From the optical data we determine the apparent electron-impact excitation cross sections for five vibrational levels of the  $C^3\Pi_u$  electronic state of  $N_2$ . Using the Franck-Condon principle we determine that in an electron beam excitation experiment the population of the first three vibrational levels (v=0,1,2) of the  $C^3\Pi_u$  electronic state is primarily due to direct excitation whereas cascade plays a significant role for populating the next two vibrational levels (v'=3,4). The relative intensities of the various vibrational bands of the second positive system observed in a dc discharge have been measured and compared with the results of the electron-beam excitation experiment.

The use of Fourier Transform spectroscopy for detecting infrared radiation emitted in an electron-beam excitation experiment has been explored. For this purpose we have constructed a special electron-beam excitation apparatus in which the radiation from the excited atoms is directed to a BOMEM Fourier-Transform Infrared Spectrometer. An InGaAs detector covers the wavelength region from 850 nm to 1.7  $\mu$ m, and an InSb detector (with liquid nitrogen cooling) is used from 1.4  $\mu$ m to 6.7  $\mu$ m. This apparatus allows us to monitor, in electron excitation experiments, many important infrared emissions which we could not detect previously by our photomultiplier optical detection system.